



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ALGEBRA.

129. Proposed by F. ANDEREGG, A. M., Professor of Mathematics, Oberlin College, Oberlin, O.

$$\left| \begin{array}{cccc} 1 - \binom{m-1}{0} - \binom{m-1}{1} \dots \binom{m-1}{m-2} \\ 1 & 1 & - \binom{m-2}{0} \dots \binom{m-2}{m-3} \\ 1 & 0 & 1 & \dots \binom{m-3}{m-4} \\ \dots & \dots & \dots & \dots \\ 1 & 0 & 0 & 1 \binom{1}{0} \\ 1 & 0 & 0 \dots 0 & 1 \end{array} \right| e = \sum_{n=\infty}^{n=1} \frac{n^m}{n!}, \text{ where } \binom{m-2}{k} = \frac{(m-2) \dots (m-k-1)}{k!}$$

Solution by the PROPOSER.

$$\text{Let } f(x) = e^{ex} = 1 + e^x + \frac{e^{2x}}{2!} + \frac{e^{3x}}{3!} + \frac{e^{4x}}{4!} + \frac{e^{5x}}{5!} + \dots + \frac{e^{rx}}{r!} + \dots$$

$$= 1 + (1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \dots)$$

$$+ \frac{1}{2!} (1 + 2x + \frac{(2x)^2}{2!} + \frac{(2x)^3}{3!} + \frac{(2x)^4}{4!} + \frac{(2x)^5}{5!} + \dots)$$

$$+ \frac{1}{3!} (1 + 3x + \frac{(3x)^2}{2!} + \frac{(3x)^3}{3!} + \dots)$$

$$\dots \dots \dots$$

$$+ \frac{1}{r!} (1 + rx + \frac{(rx)^2}{2!} + \frac{(rx)^3}{3!} + \dots)$$

$$\dots \dots \dots$$

$$\dots \dots \dots$$

$$= e + \sum_{r=1}^{r=\infty} a_r x^r, \text{ where } a_r = \frac{1}{r!} \sum_{k=1}^{k=\infty} \frac{k^r}{k!}.$$

$$\text{But } f(x) = f(0) + \frac{f'(0)}{1!} x + \frac{f''(0)}{2!} x^2 + \dots$$

$$\therefore f^m(0) = \sum_{k=1}^{k=\infty} \frac{k^m}{k!} \dots (1).$$

$$\text{Let } y = e^{ex}, \text{ then } \log y = ex.$$

$\therefore \frac{dy}{dx} = ye^x$. And by Leibnitz's Formula,

$$f^m(0) = f^{m-1}(0) + (m-1)f^{m-2}(0) + \frac{(m-1)(m-2)}{2!}f^{m-3}(0) + \dots$$

$$+ (m-1)f'(0) + f(0) = \sum_{k=1}^{k=\infty} \frac{k^m}{k!} \text{ [by (1)]} \quad (2).$$

Or, letting x_m represent $f^m(0)$, etc.,

$$x_m - x_{m-1} - C_{m-1,1}x_{m-2} - \dots - C_{m-1,1}x_1 = f(0) = e.$$

$$x_{m-1} - x_{m-2} - C_{m-2,1}x_{m-3} - \dots - C_{m-2,1}x_1 = e$$

$$\dots\dots\dots$$

$$\dots\dots\dots$$

$$x_2 - x_1 = e$$

$$x_1 = e$$

$$\therefore x_m = \begin{vmatrix} e & -\binom{m-1}{0} & -\binom{m-1}{1} & \dots\dots\dots & -\binom{m-1}{m-2} \\ e & 1 & -\binom{m-2}{0} & \dots\dots\dots & -\binom{m-2}{m-3} \\ e & 0 & 1 & \dots\dots\dots & -\binom{m-3}{m-4} \\ e & 0 & 0 & 1 & \dots\dots\dots & -\binom{m-4}{m-5} \\ \dots\dots\dots & \dots\dots\dots & \dots\dots\dots & \dots\dots\dots & \dots\dots\dots & \dots\dots\dots \\ e & 0 & 0 & 0 & \dots\dots\dots & 1 & -\binom{1}{0} \\ e & 0 & 0 & 0 & \dots\dots\dots & 0 & 1 \end{vmatrix}$$

$$\therefore \begin{vmatrix} 1 & -\binom{m-1}{0} & -\binom{m-1}{1} & \dots\dots\dots & -\binom{m-1}{m-2} \\ 1 & 1 & -\binom{m-2}{0} & \dots\dots\dots & -\binom{m-2}{m-3} \\ 1 & 0 & 1 & \dots\dots\dots & -\binom{m-3}{m-4} \\ 1 & 0 & 0 & 1 & \dots\dots\dots & -\binom{m-4}{m-5} \\ \dots\dots\dots & \dots\dots\dots & \dots\dots\dots & \dots\dots\dots & \dots\dots\dots & \dots\dots\dots \\ 1 & 0 & 0 & 0 & \dots\dots\dots & 1 & -\binom{1}{0} \\ 1 & 0 & 0 & 0 & \dots\dots\dots & 0 & 1 \end{vmatrix} = \sum_{k=1}^{k=\infty} \frac{k^m}{k!} \dots\dots (3).$$

By using (2) or (3) the following special relations have been obtained :

$$2e = \sum_{k=1}^{k=\infty} \frac{k^2}{k!}, \quad 5e = \sum_{k=1}^{k=\infty} \frac{k^3}{k!}, \quad 15e = \sum_{k=1}^{k=\infty} \frac{k^4}{k!}, \quad 52e = \sum_{k=1}^{k=\infty} \frac{k^5}{k!}, \quad 203e = \sum_{k=1}^{k=\infty} \frac{k^6}{k!}, \quad 877e = \sum_{k=1}^{k=\infty} \frac{k^7}{k!},$$

$$4140e = \sum_{k=1}^{k=\infty} \frac{k^8}{k!}, \quad 21147e = \sum_{k=1}^{k=\infty} \frac{k^9}{k!}, \quad 115975e = \sum_{k=1}^{k=\infty} \frac{k^{10}}{k!}.$$

Also solved by G. B. M. ZERR.

GEOMETRY.

160. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Let GFH be the spherical triangle formed by joining the mid-points of the sides of the spherical triangle ABC ; E the spherical excess of ABC ; β , p the base and altitude of GFH . Prove $\sin \frac{1}{2}E = \sin \beta \sin p$.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Let $BA=c$, $GF=\gamma$, $FH=\beta$, $GH=\delta$, $GP=p$.

Draw AL , BM , CK perpendicular to $DGFC$. Now $DABE = DGFE = \pi$.

$AD=BE=\frac{1}{2}(\pi-c)$, $DL=ME$, $LG=$

CK , $KF=FM$.

$\therefore 2(DL + GK + KF) = \pi$. Also $2GK + 2KF = 2\gamma$.

$\therefore 2DL + 2\gamma = \pi$, or $DL = \frac{1}{2}\pi - \gamma$.

$\angle DAL = \angle EBM$, $\angle LAG = \angle GCK$,
 $\angle KCF = \angle FBM$.

$\therefore 2\angle DAL + C + A + B = 2\pi$.

$\therefore \angle DAL = \pi - \frac{1}{2}(A + B + C) = \pi - s = \frac{1}{2}(\pi - E)$.

$\cos DAL = \sin D \cos DL$. $\cos \frac{1}{2}(\pi - E) = \sin D \cos(\frac{1}{2}\pi - \gamma)$.

$\therefore \sin \frac{1}{2}E = \sin D \sin \gamma$.


Now $\sin D : \sin DFH = \sin \beta : \sin DH$. But $DH = \frac{1}{2}\pi$.

$\therefore \sin D = \sin \beta \sin DFH$. But $\sin p = \sin \gamma \sin DFH$.

$\therefore \sin D = (\sin \beta \sin p) / \sin \gamma$.

$\therefore \sin \frac{1}{2}E = \sin \beta \sin p$.

Also solved by J. SCHEFFER and L. C. WALKER.



161. Proposed by MARCUS BAKER, U. S. Coast and Geodetic Survey Office, Washington, D. C.

A circle, radius r , is inscribed in a triangle ABC . In the angles A , B , and C are inscribed circles each touching two sides and the inscribed circle. There are six such circles. The first group of three have their centers between the in-centers and the vertices, and the second group of three does not. Let r_a , r_b , r_c denote the radii of the first group. Then this well known relation holds : $r = \sqrt{(r_a r_b)} + \sqrt{(r_b r_c)} + \sqrt{(r_c r_a)}$. Let R_a , R_b , R_c denote the radii of the second group. Then this relation holds :